

Citation for published version:

Stokes, K, Jones, B, Bennett, M, Close, G, Gill, N, Hull, J, Kasper, A, Kemp, S, Mellalieu, S, Peirce, N, Stewart, B, Wall, B, West, S & Cross, M 2020, 'Returning to Play after Prolonged Training Restrictions in Professional Collision Sports', *International Journal of Sports Medicine*, vol. 41, no. 13, pp. 895-911. <https://doi.org/10.1055/a-1180-3692>

DOI:

[10.1055/a-1180-3692](https://doi.org/10.1055/a-1180-3692)

Publication date:

2020

Document Version

Peer reviewed version

[Link to publication](#)

(c) Georg Thieme Verlag KG, 2020.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Returning to Play after Prolonged Training Restrictions in Professional Collision Sports

Keith A. Stokes ^{1,2} k.stokes@bath.ac.uk

Ben Jones ^{3,4,5,6,7} b.jones@leedsbeckett.ac.uk

Mark Bennett ^{8,9} markbennettanalytics@gmail.com

Graeme L. Close ^{10,2} G.L.Close@ljmu.ac.uk

Nicholas Gill ^{11,12} nicholas.gill@nzzrugby.co.nz

James H. Hull ¹³ J.Hull@rbht.nhs.uk

Andreas M. Kasper ^{10,2} A.Kasper@2011.ljmu.ac.uk

Simon P.T. Kemp ² simonkemp@rfu.com

Stephen D. Mellalieu ¹⁴ smellalieu@cardiffmet.ac.uk

Nicholas Peirce ¹⁵ Nick.Peirce@ecb.co.uk

Bob Stewart ² BobStewart@RFU.com

Benjamin T Wall ¹⁶ B.T.Wall@exeter.ac.uk

Stephen W. West ¹ s.west@bath.ac.uk

Matthew Cross ^{17,1} m.cross@bath.ac.uk

1 University of Bath, Bath, UK

2 Rugby Football Union, Twickenham, UK

3 Carnegie Applied Rugby Research (CARR) centre, Carnegie School of Sport, Leeds Beckett University, Leeds, LS6 3QS, United Kingdom

4 Leeds Rhinos Rugby League club, Leeds, LS5 3BW, United Kingdom

5 England Performance Unit, The Rugby Football League, Leeds, LS17 8NB, United Kingdom

6 Division of Exercise Science and Sports Medicine, Department of Human Biology, Faculty of Health Sciences, the University of Cape Town and the Sports Science Institute of South Africa, Cape Town, South Africa

7 School of Science and Technology, University of New England, Armidale, NSW, Australia.

8 Rugby Union of Russia, Moscow, Russia

9 Applied Sport Technology Exercise and Medicine Research Centre (A-STEM), College of Engineering, Swansea University, Swansea, Wales

10 Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK

34 11 New Zealand Rugby Union, 6011 Wellington, New Zealand
35 12 Te Huataki Waiora School of Health, Adams Centre, The University of Waikato, 3116
36 Tauranga, New Zealand
37 13 Department of Respiratory Medicine, Royal Brompton Hospital, London, UK
38 14 Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff,
39 UK
40 15 England and Wales Cricket Board, National Cricket Performance Centre, Loughborough
41 University, Loughborough, UK
42 16 School of Sport and Health Sciences, College of Life and Environmental Sciences,
43 University of Exeter, Exeter, UK
44 17 Premiership Rugby, Twickenham, UK
45

Returning to play after prolonged training restrictions in professional collision sports

Abstract

The COVID-19 pandemic in 2020 has resulted in widespread training disruption in many sports. Some athletes have access to facilities and equipment, whilst others have limited or no access, severely limiting their training practices. A primary concern is that the maintenance of key physical qualities (e.g., strength, power, high-speed running ability, acceleration, deceleration and change of direction), game-specific contact skills (e.g., tackling) and decision-making ability, are challenged, impacting on performance and injury risk on resumption of training and competition. In extended periods of reduced training, without targeted intervention, changes in body composition and function can be profound. However, there are strategies that can dramatically mitigate potential losses, including resistance training to failure with lighter loads, plyometric training, exposure to high-speed running to ensure appropriate hamstring conditioning, and nutritional intervention. Athletes may require psychological support given the challenges associated with isolation and a change in regular training routine. Whilst training restrictions may result in a decrease in some physical and psychological qualities, athletes can return in a positive state following an enforced period of rest and recovery. On return to training, the focus should be on progression of all aspects of training, taking into account the status of individual athletes.

Keywords

COVID-19, Rugby, coronavirus, detraining, retraining, disuse

Introduction

Collision sports such as rugby union and rugby league (i.e., rugby) have different demands than many other team sports (e.g., soccer, hockey, cricket) due to multiple contact / collision game events [1]. Athletes require well-developed specific physical qualities to perform optimally [2] and mitigate the risk of injury. These physical qualities are typically developed through well-planned, periodised training programmes [3]. The preparation, maintenance and recovery of athletes is relatively well understood within a typical season [4,5], and practitioners have a wealth of experience in supporting athletes under normal circumstances.

In 2020, the spread of a coronavirus disease (COVID-19) resulted in a world-wide pandemic. As a consequence social measures have been implemented that preclude sports competition and many aspects of team sport training. A primary concern, and the motive for this review, is that development and maintenance of key physical qualities (e.g., strength, power, high-speed running ability, acceleration, deceleration and change of direction), game-specific contact skills (e.g., tackling) and decision-making ability, is challenged during physical distancing and movement restriction measures as a consequence of COVID-19. Players are unlikely to be able to train together as teams in any form, access training facilities or public gymnasiums, nor have routine access to coaching, conditioning and medical staff. Indeed, the majority of elite athletes will be attempting to train at home within the constraints of the equipment and space that they have available to them. Some players will have access to excellent training facilities in their home, some will have access to limited facilities, and some might have no access to equipment or adequate space at all. The variation in training activities that athletes can undertake during a period of restriction will likely present additional challenges when planning the resumption of team training. As such, the specific needs of each individual athlete will require consideration upon the return of training and competition.

Although the likely impact of the COVID-19 pandemic is unprecedented in scale, there are examples of the consequences of enforced restriction of access to training on returning to sport. For example, following a 20-week lockout in the National Football League in 2011, on returning to sport there were more frequent soft tissue injuries [6,7]. Therefore, with a focus on rugby league and rugby union, the purpose of this review is to examine the available evidence related to; potential changes to physical qualities and function during the period of modified training, strategies to mitigate this decline in function, and the time taken to return

players and teams to “game ready” status. It is anticipated that many of the principles outlined in this review will be applicable to a broader range of collision sports (e.g., American football, Australian football). The final section provides practical recommendations that focus on restarting these sports after an extended break from training.

Physical qualities for rugby

The demands of rugby require athletes to have high levels of lower-body and upper-body strength and power [8]. Rugby players have high levels of lean mass [9], in comparison to other sports (e.g., soccer) [10], in addition to well-developed aerobic and anaerobic running capacities [11,12]. Strength and power are related to general athletic (e.g., speed, acceleration and change of direction) [13] and rugby specific (e.g., tackling) [14,15] qualities. The tackle and other contact events (e.g., ruck, maul, scrum) require high levels of strength and power to overcome resistant forces from opposition players.

Within rugby league and rugby union, strength and power have been shown to vary between age [16,17], and playing position [2,18]. Professional rugby league players have been shown to have greater strength and power than semi-professional or amateur players [17]. Strength appears similar for professional and semi-professional rugby union players, whereas professional players have greater power [8]. Furthermore, stronger players with higher levels of aerobic fitness have been shown to recover more quickly following rugby league match play [19].

Considerations for injury in relation to enforced modified training

There are numerous conceptual models that identify risk factors for injury (e.g., strength, training load, competition schedule, previous injury) [e.g., 20,21]. However, the evidence for proposed risk factors for injury in elite sports settings is often not as clear as might be expected, perhaps because athletes who are competing have usually reached an explicit or implicit minimum physical requirement for participation at a given level of play.

There is evidence of an association between strength of specific muscle groups and overuse shoulder injuries in throwing sports [e.g., 22] and groin injuries in a range of sports [e.g., 23]. In the case of hamstring injury risk, evidence of an association between strength and injury is

mixed (for detailed review see [24]), although a combination of performing eccentric Nordic hamstring exercises [25] and regular exposure to high speed running [26-28] appear to be protective against hamstring injuries.

A high proportion of injuries in collision sports are associated with contact mechanisms, for example, the tackle is associated with ~50% of all injuries in professional rugby union [29,30]. Even in a training environment, the greatest incidence of injury is in full contact training [31]. Therefore the ability of tissues to withstand substantial acute external forces may be key. In a prospective cohort study of professional rugby league players, those with poorly developed high-speed running ability (hazard ratio (HR): 2.9, 95% confidence interval (CI) = 1.7-4.0) and upper-body strength (HR: 2.2, 95% CI = 1.3-3.7) had a higher incidence of contact injury [32].

A systematic literature review and meta-analysis [33] identified six studies that have examined the effect of strength training interventions on injury outcomes in military [34] and elite [35,36], amateur [36,37] and youth [38,39] soccer. All of the interventions reduced injuries, with 95% certainty of more than halving injury risk (average reduction, 66%, 95% CI 52% to 76%). These findings provide compelling evidence of a role for development of strength in injury prevention, although none of the studies were in professional collision sport settings. Similarly, rugby specific injury prevention exercise programmes that focus on strength, balance and proprioception [40] substantially reduced injury and concussion incidence in cluster randomised controlled trials in youth [41] and community adult [42] cohorts. Of particular interest in the context of these studies is the potential importance of neck strength in protecting against concussion [43].

A key concern arising from a period of enforced modified training due to COVID-19 is that athletes cannot maintain physical qualities that likely protect against injury. A twenty-week shutdown of the National Football League in 2011 was associated with a four-fold increase in Achilles tendon ruptures in the first 29 days of a condensed return to competition period [6]. Over the subsequent season, soft tissue injuries (considered to be conditioning-related injuries) were higher than preceding or subsequent seasons [7]. In professional rugby union, even after a short off-season typically lasting 4-5 weeks during which athletes have opportunities to train (e.g., access to gym and other training facilities), there is a greater frequency and burden of training injuries in the early, compared with later, period of pre-

season (Figure 1). This highlights one of the challenges when athletes return following an extended period of enforced modified training.

FIGURE 1 HERE

On resumption of competition, it is possible that multiple games per week are scheduled to make up for the time lost. Limited time between matches during periods of fixture congestion has been shown to be associated with more injuries in soccer [44]. Clearly, the timing and structure of reconditioning, and fixture scheduling upon resumption of the competitive season, have the potential to impact on injury outcomes.

The concept of preparedness for training and/or competition has been investigated in professional rugby union, with intermediate cumulative load over four weeks showing a likely beneficial reduction in injury risk compared with low or high four-week loads [45]. In the same study, sudden increases (or spikes) in training load were shown to increase the risk of injury [45]. Exposure to competitive matches also appears to influence injury risk in professional rugby union, with involvement in less than 15 or more than 35 games over a 12-month period associated with a greater injury risk than being involved in between 15 and 35 games [46]. An extended period without competition will result in more players having played a low number of games in 12 months, potentially increasing injury risk.

In returning players to competition, standard considerations around individual risk factors will be important to consider. A potential positive of a period of modified training practices and no matches, is that it may allow for prolonged rest that is rarely afforded to professional rugby players. Previous injury has consistently been shown to increase subsequent injury risk [47], and there may be opportunity to focus on full recovery and rehabilitation from previous injuries, although restricted access to appropriate rehabilitation modalities might limit any positive impact. Given that some subsequent injuries considered to be, in part, related to inadequate rehabilitation [47], individual management of athletes when returning to full training is required. The Strategic Assessment of Risk and Risk Tolerance (StARRT) framework may be helpful in this respect [48]. Some athletes may even have developed injuries during the period of restricted training due to enforced changes in training type, timing, load and surface. A further consideration is that athlete anxiety may be elevated by a number of facets of an extended period of modified training due to COVID-19, which might impact on injury risk when returning to play [49].

Under normal circumstances, most elite collision sport players will be conditioned to a level that is protective against injury. However, a sustained period of enforced modified training is likely to impact upon this conditioning and is likely to result in increased injury risk. It is important to consider strategies to mitigate losses in physical function and to develop appropriate reconditioning strategies. These should be considered on an athlete-by-athlete (e.g., training status and injury history), sport-by-sport (e.g., the stage of the season), and country-by-country (e.g., local government COVID-19 guidelines) basis.

Potential changes in physiological characteristics in response to reductions in training

Athletes' musculoskeletal, respiratory and cardiovascular systems are accustomed to a large volume and intensity of training stimulus, and any considerable reduction in habitual stimuli will lead to a degree of physiological system and tissue deconditioning, in turn reducing physical performance. There is limited evidence regarding detraining in elite athletic populations, but principles of deconditioning can be translated from human laboratory studies using extreme experimental models such as limb immobilisation (local disuse), bed rest (whole body disuse) or reduced step count (moderate decreases in physical activity) in previously 'healthy' individuals [50]. Regardless of the model, such studies reliably show that deconditioning is rapid and profound [50].

Physical inactivity quickly leads to a myriad of interrelated cardiovascular deconditioning responses. Experimental bedrest [51] and short term detraining in trained individuals [52] decreases plasma volume, reduces baroreflex sensitivity, impairs the sensitivity with which the vasculature can appropriately redistribute blood volume, interferes with heart rate and blood pressure regulation, induces cardiac muscle atrophy and impairs myocardial mechanics and stroke volume. Though the time course and severity of some of these responses has not been precisely delineated, their integrated nature rapidly impairs oxygen (and nutrient) delivery and tissue extraction, and can be expected within less than four weeks [52].

Skeletal muscle appears particularly susceptible tissue to disuse-induced deconditioning, with substantial impairments in markers of metabolic health (reflecting declining muscle tissue quantity and quality) within just one week [53,54]. Disuse also almost immediately reduces daily muscle protein synthesis rates [55], largely driven by a reduced ability of the inactive tissue to extract dietary derived amino acids from the circulation following each meal [56] and utilise them for the construction of new proteins [57]. The resulting loss of muscle mass can be-nearly 100 g after only two days of limb immobilisation [55]. This increases to >250 g after one week, while one week of bed-rest results in ~1.5 kg of whole body muscle loss [53]. Strikingly, muscle strength and force generating capacity of a muscle group subjected to extreme disuse declines by ~1.5-2% per day [58], around 3-fold higher than the loss of muscle mass alone [59].

Muscles (groups) of a higher ‘training status’ *within an individual* (i.e., higher habitual gravitational loading and mechanical workload) typically decondition more rapidly. For example, the quadriceps atrophy more rapidly than the hamstrings [60] and the large postural and gross motor movement muscles of the legs, trunk and back atrophy more quickly than the arms or other smaller muscles more attuned to fine movements [61]. In the event that disuse is brought about by any type of acute injury, which would typically elicit a local and systemic inflammatory response, muscle loss may be further accelerated over rest alone [62]. At the single fibre level, muscle disuse atrophy is characterised by a decrease in cross sectional area of all muscle fibre types, though type II fibres appear particularly susceptible [63]. Skeletal muscle fibre atrophy is accompanied by considerable and disproportionately large declines in function at the level of the muscle fibre. Despite this, some evidence points towards deconditioning bringing about a ‘faster’ overall muscle phenotype, seemingly due to increased expression in the faster isoforms of muscle myosin heavy chain across fibre types [64] rather than any ‘fibre type switching’.

Bone demineralisation also occurs within a week of unloading [65], while tendon and connective tissues such as ligaments appear to be more resistant to short-term muscle disuse [66,67], likely due to considerably lower protein turnover rates. However, within a month of detraining, impairments in tendon and ligament tensile strength and functionality can be expected [68]. Deconditioning of the tendon and ligament tissue also impacts on metabolic

and functional performance [66,67]. The crucial structural role of these collagen rich tissues within the musculoskeletal system (particularly within joint stabilisation) contributes heavily to movement and force generation, and therefore their deconditioning also contributes to degenerate physical performance.

It is clear that the extreme models of disuse described above do not reflect the experiences of athletes during most periods of training restriction. However, case-study data on elite footballers suggest that injury induced periods of disuse (e.g., Anterior cruciate ligament [ACL] surgery) lead to rapid tissue and performance detriments that reflect the changes seen in laboratory trials (e.g., loss of muscle mass and function, gain in fat tissue and alterations in metabolic rate) [69,70]. Such effects are evident despite ‘best practice’ in terms of nutritional and physical therapy countermeasures being applied. Furthermore, elite athletes reducing training at the end of their competitive season can expect rapid (within 5 weeks) declines in function, with the extent being related to the level of withdrawal from training [71]. Such data brings into stark focus the challenges that those involved in collision sports face if athletes undergo enforced periods of reduced or absent training load.

Maintaining muscle mass and function during enforced modified training

Fortunately, even in extreme physiological models of disuse, small amounts of exercise can mitigate losses in muscle size and function. For example, eighty-four days of bed rest in healthy men resulted in a 17% reduction in muscle size and around 40% reduction in muscle strength and power [72]. However, when maximal concentric supine squats were performed every third day, muscle size, strength and power were maintained [72]. In 60 days of bed rest, just three minutes of “supine jumps” on 5-6 days per week maintained leg lean mass and strength, compared with reductions of around 10% without exercise [73]. These examples are important in illustrating the concept of mitigating losses in muscle mass and function during deconditioning, but cannot be translated directly into sports settings.

A key question when access to training facilities is limited is whether heavy loads during resistance training are required for the development, or maintenance, of muscle mass and strength. During resistance exercise all motor units are recruited at momentary muscular

failure, regardless of the load used [74]. In turn, rates of muscle protein synthesis for up to 24 hours after exercise were similar when healthy men performed knee extension at 30% of one repetition maximum (1RM) to failure compared with 90% 1RM to failure [75]. Taking this further, 10 weeks of knee-extension training to failure at 30% 1RM and 80% 1RM in healthy young men resulted in similar change in quadriceps volume (hypertrophy), although gains in strength as assessed by 1RM was significantly higher following training at 80% 1RM [76]. Other studies have also reported similar hypertrophy response in lower-load and higher-load resistance training, with smaller gains in strength in lower-load training [77,78]. Furthermore, 12 weeks of whole-body resistance training at either 30-50% 1RM or 75-90% 1RM in trained individuals resulted in similar changes in whole body lean mass [79]. However, in this study, 1RM strength was tested every third week, essentially allowing a small amount of high load training in both groups, and the strength outcomes were similar in all tests other than bench press for which there was a small but significantly superior gain in the 75-90% 1RM group. Incorporating plyometric training might also be beneficial, given that eccentric muscle actions have the potential to induce neural adaptations, even in the absence of heavy loads, and that both concentric and eccentric peak torque were better maintained during detraining following coupled concentric and eccentric resistance training than concentric training alone [80]. Furthermore, although evidence is mixed, meta-analysis showed small-to-moderate effects of plyometric training on maximal strength in healthy adults [81].

Focusing on elite athletes, bench press and bench pull performance were assessed in kayakers before and after five weeks of detraining following the World Championships [71]. Seven athletes discontinued all training, while seven completed a dramatically reduced volume of training that included one resistance training session per week. In those that discontinued training, bench press 1RM declined by 8.9% and bench pull by 7.8%, whereas in those completing one resistance training session per week, declines in strength were much smaller at 3.9% for bench press and 3.4% for bench pull. In addition, those that discontinued training suffered a reduction in $\text{VO}_{2\text{max}}$ of 11.3%, whereas those that reduced training volume to just two endurance session per week experienced reductions of 5.6%. As a note of caution, in athletes for whom strength and power are key attributes, the possible interference effect of endurance training on strength adaptations should be considered [82,83]. This might be particularly relevant when running and cycling activities are possible, but access to resistance training facilities are limited.

From both a performance and injury prevention perspective, incorporating high speed running into training is likely to be beneficial. Sprint training has been demonstrated to have positive impacts on hamstring architecture and sprint performance [84], and regular exposure to maximal running velocity has the potential to reduce injury risk [26]. The addition of both eccentric hamstring training [25] and plyometric training [85] may also be appropriate.

Practically, strength and power trained athletes may find it difficult to match the loads needed to maintain size, strength and power. Performing resistance training to momentary failure, even with low loads, may mitigate some losses in muscle size, and if some training with high resistance can be incorporated, even if not at the usual frequency, it is possible to maintain strength characteristics. Alternatively, plyometric exercises might provide a sufficient neural stimulus to contribute to the maintenance of strength. Furthermore, given that neural adaptations might be retained for longer than 12 weeks in trained individuals [64], and that, even if this is not the case, neural adaptations occur early in response to resistance training [86], a focus on retaining as much muscle mass as possible during restricted training is recommended, followed by the re-introduction of high resistance in training once access to facilities and support is possible.

Psychological considerations during enforced modified training

The training limitations arising from COVID-19 present a number of psychological considerations which may influence preparation for, and subsequent return to, rugby competition. These include the impact of confinement and isolation, deconditioning effects, deterioration in skill execution/performance, and, the opportunity for recovery and posttraumatic growth.

In addition to the psychological effects from periods of confinement and isolation reported in the general public [87], such as post-traumatic stress symptoms (i.e., depression, anxiety, confusion, and anger), athletes may be at further risk due to the impact on their athletic identity. Athletic identity refers to the extent to which an individual identifies with their role as an athlete [88]. Any challenges to the ability to reinforce this identity through reduced capacity to train, play and achieve goals (typically seen in injured or retired athletes) are associated with feelings of loss, identity crisis and distress [89]. While engaging with social

support networks is seen as a key resource to cope with potential threats to athletic identity arising from the restrictions, it is likely athletes will be socially isolated from those who contribute most to supporting their sense of athletic identity (teammates, staff, fan base). An extended period of isolation from fellow team mates is also likely to impact upon the social and psychological group process that underpin a team's effectiveness to work together (i.e., teamwork; [90]) and subsequently perform.

In contrast to the physiology literature, limited research has examined the psychological effects of a period of detraining or rest. While acute bouts of rest (e.g., 2-week mid-season break) improve subjective perceptions of some aspects of wellness, such as fatigue and muscle soreness [91], there is no evidence examining the chronic effects of deconditioning. In the professional practice literature, Bompa and Buzzichelli [92] suggest an abrupt cessation of training by highly trained athletes creates a phenomenon known as detraining syndrome, characterized by insomnia, anxiety, depression, alterations to cardiovascular function, and loss of appetite. These symptoms are usually not deemed pathological and can be reversed if training is resumed within a short time, however, with prolonged cessation, symptoms may become more pronounced.

The principle of reversibility dictates that athletes lose the beneficial effects of training on cessation/reduction in such activities [93]. A decline in skill execution/performance may therefore be expected from a lack of deliberate team or individual skill-based practice, and will vary with the nature and type of skill [94]. Offsetting skill reversibility will rely in part on the ability to assess the relevant elements of the required skill performances, and utility of the practice-based knowledge regarding retention or transfer effects that accompany practice of these skills [95]. The use of the cognitive technique of imagery, specifically mental rehearsal of the execution of individual skills/team strategies, can aid with physical skill learning or refinement [96]. However, no research has considered the role of imagery in skill retention following deconditioning or rest. Video-based observation (modelling) of existing skill execution or performance can also be used to promote physical skill learning and refinement, and can enhance both individual and teams confidence in their ability to execute the skill [97].

A period of abstinence from sport may also offer athletes an opportunity for mental rest and recovery, especially where restrictions occur towards the end of a competitive season. Recent

research in professional rugby union [98] suggests advanced information regarding the timing and length of any competition break (i.e., off-season) can determine the level of autonomy players perceive over their break from the sport, and the subsequent degree of psychological recovery achieved. Given restrictions associated with the COVID-19 pandemic have meant a suspension (as opposed to termination) in the current competition season, athletes are being asked to engage in a level of interim individual training that does not align to a designated off-, pre- or in-season period, without any competition goal or outcome to pursue. This training ‘limbo’ may reduce players’ ability to cognitively ‘detach’ [99] and negate any potential psychological benefit associated with time away from the sport.

In considering the human trauma associated with COVID-19 it is noteworthy that the consequences for mental health and wellbeing will not be inherently negative. Potential exists for growth in response to traumatic life experiences, where growth involves profound and transformative positive changes in cognitive and emotional life that are likely to have behavioural implications [100]. Research in sport has examined growth in relation to adverse intrapersonal experiences such as long term injury and sport retirement [101], and recently at the interpersonal and organizational level (see [102]). Both individual and collective psychological growth may be derived from the trauma and adversity athletes, teams and their staff face during the restrictions. The extent to which growth is likely occur will, however, be influenced by the amount and nature of the support provided before, during, and after the restrictions.

Nutritional considerations during enforced modified training and re-training

The overarching goal during a phase of restricted training is to maintain physical capacity via preservation of muscle mass, minimisation of unwanted body fat increase, support of immune function and maintenance of cardiovascular capacity. Energy expenditure may be reduced during a period of reduced training, although other factors may be increased contributing to overall energy expenditure. For example, Anderson et al. [103-105] suggests that with injured athletes who have a reduction in their absolute training intensity, increases in other factors (e.g., frequency of resistance training and rehabilitation) result in trivial changes in total daily energy expenditure (estimated reduction of 300 kcal·d⁻¹). Therefore nutritionists should consider an individual’s habitual physical activity level (e.g., dog walking, living and training

logistics, active family) prior to suggesting a reduced total caloric intake. One of the main challenges for bespoke nutritional intervention during this period will be the accurate assessment of daily energy expenditure with a '*one-size fits all*' approach being particularly problematic. Rugby players have large inter-individual differences in daily energy expenditure when measured via doubly labelled water, even when the players appear to be undertaking similar training sessions (Table 1) [106-109]. This highlights the substantial contribution of activities away from the training ground on total daily energy expenditure and it is therefore essential that nutritionists attempt in some way to quantify the activities of the day during this period of training restriction.

TABLE 1 HERE

Research has shown decreased insulin sensitivity, attenuation of postprandial lipid metabolism, and an increase in fat mass as a consequence of simply reducing step count (~1300 from ~10000) for 2-3 weeks [110] alongside increases in visceral adiposity [111]. If athletes reduce their daily activities, there is a requirement to reduce caloric intake versus 'normal' habitual competition, however it is important to maintain habitual protein intake. Although the majority of research has focused on middle- and older-aged males [112,113], targeted nutrition, specifically dietary protein intake, has been shown to mitigate the consequences of reduced activity, even in younger adults [114,115]. One specific essential amino acid that may play the most pivotal role in the attenuation of anabolic resistance as a result of disuse is leucine, a potent stimulator of mTOR and thus muscle protein synthesis [116]. It is therefore suggested that athletes maintain a high protein diet rich in leucine, consuming approximately 0.4 g·kg⁻¹ body of protein regularly (every 4 hours) throughout the day [117]. The reduction in calories will therefore come from reduced carbohydrate and fat intake utilising a periodized carbohydrate model based on the demands of the training day [118]. It is important however that sufficient carbohydrates are consumed during this period and that athletes do not adopt a ketogenic style diet given the strong links between carbohydrates, stress hormone responses and the immune function (discussed further in [119-121]). Protein is often used in conjunction with creatine monohydrate to support maintenance/gains in strength and lean mass. Supplementation has been shown to attenuate loss of upper arm mass and strength, specifically during times of disuse (limb

immobilisation), as well as increase muscle hypertrophy following lower limb immobilisation [122,123].

From an immune support perspective, research has shown that protein may also have a pivotal role in supporting the immune function, specifically antibody response to infection [124,125] again highlighting the need to maintain sufficient protein intakes. Other nutritional factors that may aid with microbe ‘resistance’/‘tolerance’ during this specific period include supplementation of 500-1000 mg vitamin C [126], 1000-4000 iU daily vitamin D₃ [127,128] and ~20 billion CFU multistrain probiotic [120,129-131]. For a full review of nutrition and immune tolerance the reader is referred to Walsh [120].

Reconditioning considerations on return to training

Extended periods of restricted or modified training create a challenge for athletes when returning to sport ready to perform and with a low risk of injury. Following the National Football League “lockout” in 2011 it is not known whether the athletes returned in good physical condition or not, but the increased incidence of Achilles tendon injuries [6] suggests that athletes may not have been physically ready for the demands of the game or the return to play protocols were not thorough and progressive enough.

Some physical qualities are likely easier to maintain (e.g., strength, power, aerobic and anaerobic capacity and linear speed) with minimal equipment, although on return to training, all require consideration. In many cases players have to train alone without access to equipment, appropriate space or expertise, leading to an inability to maintain the required intensity of training. This will vary between countries, given variations in government-enforced physical distancing protocols. The most difficult aspect of rugby training to replicate when training individually are the “intricacies” of the sport. These include the sport-specific physical and mental demands, such as changes in direction while running at speed, running with ball in hand, attempting to evade would-be tacklers and then being tackled, lineout jumping, cutting, tackling, scrummaging, ruck clearance and mauling [132]. In this context, decision-making can only be practiced when training with others. Typically, athletes would return to structured preparation after a 3- to 6-week off-season and progress to playing the game over 6 to 12 weeks.

496

497 It is vital that athletes returning to rugby following a period away from team training
498 undertake a well-planned, progressive return to play programme to prepare to perform and to
499 decrease the risk of injury (see [133]). High-speed (or sprinting) running is one specific
500 consideration on return to training, given the concurrent benefit to performance (e.g.,
501 acceleration and maximum sprint speed [134]) and injury prevention [135]. High-speed
502 running exposure should be managed carefully as an excess or rapid increase in training load
503 may increase soft tissue injury risk [21]. In sprinters, the training phase (e.g., following the
504 off-season) and transition phase between the preparation period and competitive season
505 appear to be vulnerable periods for injury [136]. That said, high-speed running is paramount
506 for sprint performance enhancement [137] and the morphological and architectural lower
507 limb qualities [84] suggesting it should be incorporated into reconditioning training
508 programmes. On return to match play, if the difference between training speed and
509 competition speed is large, this may also increase injury risk [85], although empirical
510 evidence does not exist to support this.

511

512 For athletes that have had limited or reduced exposure to high-speed running, the initial
513 weeks of training should focus on the re-familiarisation of the intensity and duration required
514 for training and competition, which should be progressed gradually [85]. No clear
515 recommendations exist on sprint exposure for rugby players, although general principles such
516 as avoiding high-speed running on consecutive days do exist [85]. These should be
517 considered alongside other training modalities rugby players are exposed to [138] and their
518 potential interaction (e.g., avoiding high velocity sprinting following fatiguing lower-body
519 resistance training). As a guide, athletes should be exposed to a range of sprint distances, to
520 allow the development of acceleration (10-50 m, >98% intensity, total session volume 100-
521 300 m), maximal velocity from a flying start (10-30 m and >98% intensity, total session
522 volume 50-150 m) and sprint-specific endurance (80-150 m and >95% intensity, total session
523 volume 300-900 m) (see [85]). Resisted sled sprint training may also be beneficial (see
524 [139]). Given high-speed running exposure will be one of a number of qualities practitioners
525 will aim to retrain, it may be more prudent to focus of the quality of the high-speed running
526 exposure, as opposed to volume *per se*. For example, practitioners should end a high-speed
527 running session when there is a drop off in performance, and/or technical error is observed,
528 and 1-2 minutes of recovery can be provided for every second spent maximal sprinting
529 between repetitions [85,140].

530

531 Preparing for the rugby specific actions is also a key consideration for athletes and
532 practitioners. This is best achieved through performing such actions during ‘practice’
533 involving the performance of the fundamentals of the game with teammates in either
534 ‘opposed’ or ‘unopposed’ situations. For example, simple skills such as catch and pass, game
535 plan understanding, tackle progressions, and the changes in direction that occur during
536 normal practice are part of preparing to play the game.

537

538 Another example of how to integrate injury prevention and progressive exposure to game
539 play is change of direction and agility. These are important facets to evade tacklers and create
540 an open field of play [141]. An athlete's agility performance is strongly influenced by the
541 ability to rapidly decelerate and reaccelerate while adjusting his or her momentum to either
542 pursue or elude opponents [142]. While athletes could be working on change of direction,
543 acceleration, deceleration and agility by themselves (e.g., practice sharp changes in direction
544 while running at high speeds, and including rapid acceleration and deceleration), once they
545 return to training with team members and return to play progressions, the key is to gradually
546 build in layers of intensity (e.g., speed of run and sharpness of direction change) and decision
547 making.

548

549 As described previously, the tackle poses the highest injury risk in rugby, accounting for
550 around 50% of injuries [29,30]. Therefore it is important that athletes have the required
551 physical and technical skill set to perform safely and effectively. The development of specific
552 tackle skills have received little attention within the scientific literature [143], but poor tackle
553 technique has been shown to result in a higher injury risk [144,145], and fatigue has been
554 shown to alter tackle technique [146,147]. Several frameworks have been proposed on how to
555 train the tackle [148,149], although the effectiveness of these is yet to be determined. It is
556 likely that following a prolonged period of non-contact training, due to the enforced physical
557 distancing players will require a graded exposure to both the technical and physical
558 components of the tackle [148]. Following a typical 3-6-week off-season and 6-12 weeks pre-
559 season, athletes will start to engage in contact and tackle training during week 3-6, with
560 progressions over 2-6 weeks. It is likely athletes will need at least 3-4 weeks of progressions
561 and exposure to tackle and contact skill training to prepare for matches.

562

The simplest way to prepare for the explosive demands of the game is to ensure all activities follow well-planned progressions (Figure 2). Such progressions are dependent on the sport specific task in question and the position demands for each individual. In the specific context of return to training in relation to COVID-19, local government policy and risk assessments based on potential for COVID-19 transmission in any given activity or session will impact upon decisions regarding the choice and rate of progressions.

FIGURE 2 HERE

Development of strength and power on return to training

Rates of change in power and strength are influenced by the intensity (percentage of maximal), volume (sets x repetitions) and frequency of resistance training, with relatively small changes in maximal strength and power in elite athletes, due to their previous training status [3] (Tables 2 and 3). In a meta-analysis, maximal strength was reported to increase at a rate of 1.8% weekly [3]. Similarly, Issurin [150] reported that elite kayakers improved their maximal strength by 5.9% over the first 3-weeks of a 20-week training cycle, an average improvement of 1.93% a week. However, during the second 3-week week phase of training average change in strength was only 0.53% a week, and continuation of the programme resulted in minimal improvements in maximal strength, with the final 14-weeks resulting in a further total increase of only 1.82%. These results suggest that only the first 6-weeks of a strength training cycle provides positive adaptations for elite athletes. The changes in strength and power during a professional rugby season also demonstrated that the majority of strength changes occur early in a programme [5]. Improvements in strength during the first 12-weeks of training were $2.7 \pm 1.1\%$. During the second 21 weeks of training, strength gains were on average $1.9 \pm 1.1\%$. These changes are considerably lower than reported by McMaster et al. [3], however these results reflect changes in force production during an isometric squat rather than specific lifts (e.g., squat) that also improve due to familiarity of the exercise task, and technique changes. It is worthwhile noting that the greatest changes in strength in professional rugby players in England coincided with the highest volumes of strength training, during the second phase average strength loads were 63% of those utilised in the first phase [5].

TABLE 2 HERE

TABLE 3 HERE

The potential rate for gains in power for athletes in collision sports appear relatively low, however, the protocols utilised in studies examining power changes are more suited to induce changes in maximal strength than power [151,152]. Over a season of professional rugby, the greatest improvements in power were observed in the early competition phase, when strength training frequency decreased, training intensity remained high, and total volume load was reduced [5].

Rugby players are typically assessed for strength and power at various stages within the season. Whilst published data are not available, unpublished data (Bennett, Unpublished; Table 4) demonstrate the changes in strength and power exercises from 26 male Tier 1 International rugby union players over a 5-week physical training phase after a 5-week break from organised strength and power training (2-week end-of-season tour and 3-week recovery). These data provide a reference point for expected strength and power development rates following extended breaks in training, such as that resulting from the COVID-19 pandemic.

TABLE 4 HERE

Table 5 presents the changes in a male professional rugby union team (35 players) in the first 4-week training block, after a 4-week off-season (Bennett, Unpublished). Of note in both the data on the professional players and also the international players, considerably greater changes in lower body strength are observed in comparison to upper body strength in both instances. This could be a similar response to the muscle-specific atrophy described in response to extreme models of disuse [60,61]. Alternatively, it could be related to players favoured training options when away from an organised environment (e.g., undertaking unsupervised upper-body, as opposed to lower-body resistance training).

TABLE 5 HERE

Neural adaptations appear to provide a greater contribution to strength increases than muscular hypertrophy early in training [86], but changes in power and maximal strength that occur from detraining are a likely a result of both neural adaptations and a decrease in cross-sectional area of the muscle [153]. That said, there is evidence to suggest that neural changes from heavy strength training are long lasting and can extend beyond 12-weeks of detraining [64]. These findings suggest that on return to club training, hypertrophy of muscle fibres should be the primary focus, especially in those players who have lost significant muscle mass. Some evidence is present in the literature with regard to “muscle memory” a phenomena where previously trained musculature retains a considerable proportion of relevant adaptations and does not return to its pre-trained state, even after a considerable period of detraining (for review see [154]). It has been shown that individuals with a substantial strength training background can regain previous muscle fibre hypertrophy and strength levels in a relatively short period of time, as much as 32-weeks of detraining can be reversed with 6-weeks of strength and power training [155]. This, alongside a maximal window of 6-weeks before the rate of return on strength training is minimised [150], would suggest a 6-week training block is sufficient for professional rugby players to regain previous physiological adaptations.

Considerations for athletes returning to training after suspected or confirmed COVID-19 infection

Any discussion or guidance regarding re-conditioning in athletes needs to acknowledge and reflect the general principles informing return to play after acute medical illness. This is particularly important for athletes with confirmed or suspected COVID-19 infection. In many cases, an athlete will only have been given a presumptive diagnosis, based on the presence of typical clinical features (e.g., dry persistent cough and febrile illness) leading to a 7- or 14-day period of self-isolation. Many suspected cases will not have undergone formal testing due to local testing procedures and policies. Indeed, for most young, fit individuals, acute COVID-19 infection is associated with very few overt systemic features, typically only very mild upper airway symptoms (e.g., anosmia), and the athlete may often not feel unwell. A very small number of previously fit young people will develop moderate to severe disease and may require acute medical care, including in some cases, the provision of hospital-level support, and possibly ventilatory support [156]. In this latter group, data series indicate an

almost ubiquitous presence of pulmonary infiltrate (on either a chest x-ray [CXR] or computerised tomography [CT] scan) and a high prevalence (8-28%) of elevated markers of cardiac dysfunction (e.g., troponin rise) that may manifest acutely as myocarditis, heart failure, cardiac arrhythmias and acute coronary syndrome [157,158]. There also appears to be an increased risk of thromboembolic events, which need to be considered in the differential diagnosis in any clinical presentations encountered in athletes recovering from COVID-19 infection; i.e., consider deep vein thrombosis in an athlete reporting calf pain.

Historically, the most widely adopted return to play approach in athletes recovering from respiratory tract infection, is based on the ‘neck check’ approach [159]. Using this approach, athletes are advised that they may continue to exercise if their symptoms and clinical signs are confined to the upper airway (e.g., only coryzal symptoms) and a short sub-maximal exercise trial does not exacerbate symptoms. The scientific basis for this recommendation is weak, and there is long-standing concern of the potential risk of athletes with respiratory tract infection developing other clinically significant end-organ complications on their return to vigorous exercise. Of these risks, the most important is the risk of myocarditis or myocardial damage, which could be highly relevant in relation to COVID-19. The current COVID-19 pandemic, particularly challenges the ‘neck check’ approach, in that there is reported variability and an almost ‘biphasic’ recovery pattern, such that infected individuals can appear to transiently improve, only to deteriorate at a later stage; approximately one week after the onset of symptoms. In addition, and as outlined above, there is concern from emerging data, that myocardial irritation and frank myocarditis may be both prevalent and an important manifestation of this novel infection [157,158,160]. It is not yet clear if this is the case in those with clinically mild disease (i.e., in those not hospitalised), however, given the considerable cardiovascular challenge of participating in elite sport, consideration of this risk should form a key part of an individual’s return to play assessment. It is with these considerations in mind, that clinicians generally adopt a more conservative approach in planning a post COVID-19 return to play strategy for confirmed and suspected cases at the current time. Expert groups (e.g., in cardiology and respiratory medicine specialities) are starting to provide guidance for specific follow-up based on small data series of the general population and expert opinion and this will undoubtedly evolve as peer-reviewed data from the athletic population becomes increasingly available.

It is recommended that medical practitioners such as Sports Physicians, overseeing the return to training, should consider utilising an approach that incorporates and considers ‘risk’ stratification. It may also be possible to assess physiological markers including resting, exercising and recovery heart rates, beat to beat variability, ratings of perceived exertion and other indicators of reduced cardiopulmonary function. In addition, ongoing understanding of the condition may point to other markers of wider organ involvement that form part of the elite sport training monitoring such as exaggerated rises in blood creatine kinase [161] and lactate concentrations. Furthermore, a graded return to activity, perhaps akin to that used in under-recovery unexplained-under-performance syndrome [162] could be employed to guide a careful progression, whilst our understanding of the most appropriate post-COVID progression develops. In the meantime, clinicians can use Figure 3 to help inform return to play risk stratification.

FIGURE 3 HERE

Considerations for at risk groups during enforced modified training and re-training

As a result of the extended period of training restriction, there will be some athletes who are at a significantly higher risk of injury when they return to training. Although specific evidence in this area is limited due to the uncommon nature of such a period of restriction in elite sports, broader evidence available concerning predisposition for injury may assist in the identification of these at-risk groupings. For example, evidence has shown previous injury to be a strong risk factor for further injury [47]. This is particularly important to consider when the ability to a) rehabilitate and pro-actively manage any existing injuries and b) continue prehabilitation programmes for injury prevention is reduced during restriction. It is also noteworthy that following the National Football League lockdown in 2011, the Achilles tendon injured group in the early phase of return to competition were, on average, younger and had a lower exposure to the NFL environment than Achilles tendon injured players in other years, suggesting specific risk [6]. Alongside the physical health of the athlete, their mental wellbeing may also be affected, highlighting the need for well-defined and accessible support structures for athletes and staff both during and after isolation. Furthermore, and as a direct result of this extraordinary time, the best practice management of athletes who either present with COVID-19 symptoms or are returning to activity following a suspected or confirmed

case of COVID-19 is clearly of huge importance. Of note is the risk of long-term effects on the respiratory and cardiovascular systems if these individuals are not managed correctly.

Figure 4 summarises those groups considered ‘at-risk’. It is recommended that athletes that fall into these groups are given careful consideration when planning their reintegration into normal training practice. It might be suitable to utilise physical and psychological screening tools to establish a baseline upon return to the club environment and to provide practitioners with information upon which to base their periodisation and programming. Overall, an individualised approach to the at-risk groups is recommended.

FIGURE 4 HERE

Challenges and practical recommendations for collision sports

The COVID-19 pandemic has created a unique scenario for all major sports with respect to the highly unusual period of training restriction. All sporting National Governing Bodies and competition organisers will need to consider how they plan the return of training activities, and ultimately competition, balancing a range of drivers to restart sport as quickly as possible with how they best manage the welfare of their athletes. These will differ between countries (e.g., England versus New Zealand) and sports (e.g., rugby league versus rugby union) given the varying level of impact COVID-19 has had on training restrictions and modification, and the varying stages of the season athletes were in. In collision sports, the resumption of training following a period of modified isolated training will arguably be harder to manage than other sports. This is due to a number of factors which include the high-risk nature of participation and the importance of strength and power, which may be affected by restricted access to training equipment and space. In addition, the importance of skill execution in high risk areas of the game, such as the tackle, and the lack of opportunity available to train these skills during a period of restriction also requires special consideration. Even on the resumption of training, factors such as limits on the number of players that can train together and limits on the amount of time it is acceptable for players to be in close contact with on another players will influence possible training progressions. That said, the unprecedented period of non-contact training may provide a positive period for physical and psychological rest and recovery. With the application of appropriate and progressive reconditioning practices on return to training, this may improve an athletes’ performance and wellbeing.

Athletes may also be afforded the opportunity to target the development of specific physical weaknesses, without the challenges of preparing for weekly competitive matches.

Monitoring of athletes' training during the period of training restriction may be beneficial when making decisions regarding initial load and progressions when group-based training resumes. Player load monitoring should be appropriate to capture the range of stresses (e.g., volume, intensity, resistance training, running) athletes have been exposed to [163]. Microtechnology is commonly used within rugby to collect objective external load measures, but access to both hardware and software is likely to be limited when training away from club environments. Session rating of perceived exertion (sRPE; [164]) offers a practical method of monitoring player load, regardless of the exercise modality. Remote monitoring of sRPE has been shown to be valid in comparison to recall with 30 minutes of exercise cessation when collected 24 to 48 hours [165] following an activity, but not at 72 hours [166] or when collected as part of a weekly self-reported training load diary [167]. As such, athletes should aim to report their sRPE at least every 48 hrs. In addition, it might be prudent to capture information about exposure to specific training, such as high speed running. In the absence of regular monitoring during the period of training restriction, screening prior to the resumption of group-based training should capture information about the training that has been carried out by each individual athlete.

It is also logical to think that the risk of infectious transmission in contact sports is higher than in non-contact sports and so the development of medical policy to mitigate the risk of transmission alongside suspected case management is critical. Furthermore, there will be a need to assess the risk of COVID-19 transmission in close contact elements of training, and to introduce these in a graded fashion that minimises risk. Table 6 summarises the focus areas, challenges and practical recommendations that have been identified in this review that the teams and major stakeholders of elite collision sports need to consider when managing athletes during this unprecedented period of restriction and when planning the resumption of training and competition.

TABLE 6 HERE

Conclusion

The COVID-19 pandemic has created unprecedented challenges in sport, resulting in restrictions to competition and many aspects of training. These restrictions have led to concerns about the ability of athletes in collision sports to maintain key physical attributes (e.g., strength, power, high-speed running ability, acceleration, deceleration and change of direction), game-specific contact skills (e.g., tackling) and decision-making ability. Any decay in these attributes has the potential to impact on performance and injury risk on resumption of training and competition. However, with appropriate management it is possible to benefit from a rare opportunity for extended recovery and for athletes to maintain and even develop many aspects of physical and psychological function. In contrast, some physical, psychological and sport-specific attributes are challenging to affect during periods when athletes are only able to train on their own. Fortunately, a period of around 6 weeks of preparation is likely to be sufficient for most athletes to return to being competition ready, although this will ultimately depend on the length of governmental social distancing policies, which differ by country. Returning athletes to competition-ready status will require the application of broad principles of progression with the added dimension of assessing the risk of infection transmission in group training activities. Individual player circumstances should be considered from a performance and welfare perspective, particularly in the case of those athletes considered to be at higher risk of poor performance or injury.

Conflicts of Interest

No funding was received for the preparation of this review. KAS, GLC, AMK, SPTK and BS are employed by the Rugby Football Union. BJ is employed by the Rugby Football League. MB is employed by the Rugby Union of Russia. NG is employed by New Zealand Rugby Union. MC is employed by Premiership Rugby. The review was prepared based on the ethical standards of the International Journal of Sports Medicine [168]

References

- ¹ *Whitehead S, Till K, Weaving D, Jones B.* The use of micro-technology to quantify the peak match-demands of the football codes: a systematic review. *Sports Med* 2018; 48: 2549-2575
- ² *Till K, Scantlebury S, Jones B.* Anthropometric and Physical Qualities of Elite Male Youth Rugby League Players. *Sports Med* 2017; 47: 2171-2186
- ³ *McMaster DT, Gill N, Cronin J, McGuigan M.* The development, retention and decay rates of strength and power in elite rugby union, rugby league and American football. *Sports Med* 2013; 43: 367-384
- ⁴ *Argus CK, Gill ND, Keogh JW et al.* Changes in strength, power, and steroid hormones during a professional rugby union competition. *J Strength Cond Res* 2009 23; 1583–1592
- ⁵ *Gannon EA, Stokes KA, Trewartha G.* Strength and power development in professional rugby union players over a training and playing season. *Int J Sports Physiol Perform* 2016; 11: 381-387
- ⁶ *Myer GD, Faigenbaum AD, Cherny CE et al.* Did the NFL Lockout expose the Achilles heel of competitive sports? *J Orthop Sports Phys Ther* 2011; 41: 702-705
- ⁷ *Binney ZO, Hammond KE, Klein M, Goodman M, Janssens ACJW.* NFL Injuries Before and After the 2011 Collective Bargaining Agreement (CBA) arXiv:1805.01271v1 [stat.AP] 3 May 2018
- ⁸ *Argus CK, Gill ND, Keogh JW.* Characterization of the differences in strength and power between different levels of competition in rugby union athletes. *J Strength Cond Res* 2012; 26: 2698-2704
- ⁹ *Geeson-Brown T, Jones B, Till K et al.* Body composition differences by age and playing standard in male rugby union and rugby league: A systematic review and meta-analysis. *J Sports Sci* 2020; Epub Ahead of print
- ¹⁰ *Milsom J, Naughton R, O'Boyle A et al.* Body composition assessment of English Premier League soccer players: a comparative DXA analysis of first team, U21 and U18 squads. *J Sports Sci* 2015; 33: 1799-1806
- ¹¹ *Darrall-Jones J, Jones B, Till K.* Anthropometric, sprint and running profiles of English academy rugby union players by position. *J Strength Cond Res* 2016a; 30: 1348-1358
- ¹² *Darrall-Jones J, Till K, Roe G et al.* The effect of body mass on 30:15 end stage running speed in rugby union players. *Int J Sports Physiol Perform* 2016b; 11: 400-403
- ¹³ *Suchomel TJ, Nimphius S, Stone MH.* The Importance of Muscular Strength in Athletic Performance. *Sports Med* 2016; 46: 1419-49
- ¹⁴ *Speranza MJ, Gabbett TJ, Johnston RD, Sheppard JM.* Muscular Strength and power correlates of tackling ability in semiprofessional rugby league players. *J Strength Cond Res* 2015; 29: 2071-2078

- ¹⁵ *Speranza MJ, Gabbett TJ, Johnston RD, Sheppard JM*. Effect of strength and power training on tackling ability in semiprofessional rugby league players. *J Strength Cond Res*. 2016; 30: 336-343
- ¹⁶ *Darrall-Jones JD, Jones B, Till K*. Anthropometric and physical profiles of English academy rugby union players. *J Strength Cond Res* 2015; 29: 2086-2096
- ¹⁷ *Till K, Jones B, Geeson-Brown T*. Do physical qualities influence the attainment of professional status within elite 16-19 year old rugby league players? *J Sci Med Sport* 2016; 19: 585-589
- ¹⁸ *Smart DJ, Hopkins WG, Gill ND*. Differences and changes in the physical characteristics of professional and amateur rugby union players. *J Strength Cond Res* 2013; 27: 3033-3044
- ¹⁹ *Johnston RD, Weaving D, Hulin BT et al*. Peak movement and collision demands of professional rugby league competition. *J Sports Sci* 2019; 37: 2144-2151
- ²⁰ *Meeuwisse WH, Tyreman H, Hagel B, Emery C*. A dynamic model of etiology in sport injury: the recursive nature of risk and causation. *Clin J Sport Med* 2007; 17: 215-219
- ²¹ *Windt J, Gabbett TJ*. How do training and competition workloads relate to injury? The workload-injury aetiology model. *Br J Sports Med* 2017; 51: 428–35
- ²² *Achenbach L, Laver L, Walter SS et al*. Decreased external rotation strength is a risk factor for overuse shoulder injury in youth elite handball athletes. *Knee Surg Sports Traumatol Arthrosc* 2020; 28: 1202-1211
- ²³ *Whittaker JL, Small C, Maffey L, Emery CA*. Risk factors for groin injury in sport: an updated systematic review. *Br J Sports Med* 2015; 49: 803-809
- ²⁴ *Pizzari T, Green B, van Dyk, N* Extrinsic and Intrinsic Risk Factors Associated with Hamstring Injury. In: Thorborg K, Opar D, Shield A (eds.). *Prevention and Rehabilitation of Hamstring Injuries*. doi:10.1007/978-3-030-31638-9, 2020
- ²⁵ *van Dyk N, Behan FP, Whiteley R*. Including the Nordic hamstring exercise in injury prevention programmes halves the rate of hamstring injuries: a systematic review and meta-analysis of 8459 athletes. *Br J Sports Med* 2019; 53: 1362–1370.
- ²⁶ *Malone S, Roe M, Doran D et al*. High chronic training loads and exposure to bouts of maximal velocity running reduce injury risk in elite Gaelic football . *J Sci Med Sport* 2017; 20: 250-254
- ²⁷ *Malone S, Owen A, Mendes B et al*. High-speed running and sprinting as an injury risk factor in soccer: Can well-developed physical qualities reduce the risk? *J Sci Med Sport* 2018; 21: 257-262
- ²⁸ *Ruddy JD, Pollard CW, Timmins RG et al*. Running exposure is associated with the risk of hamstring strain injury in elite Australian footballers. *Br J Sports Med* 2018; 52: 919-928

- ²⁹ Williams S, Trewartha G, Kemp S, Stokes K. A meta-analysis of injuries in senior men's professional Rugby Union. *Sports Med* 2013; 43: 1043-1055
- ³⁰ Stokes KA, Locke D, Roberts S, et al. Does reducing the height of the tackle through law change in elite men's rugby union (The Championship, England) reduce the incidence of concussion? A controlled study in 126 games. *Br J Sports Med* 2019; Epub ahead of print <http://dx.doi.org/10.1136/bjsports-2019-101557>
- ³¹ West SW, Williams S, Kemp SPT et al. Patterns of training volume and injury risk in elite rugby union: An analysis of 1.5 million hours of training exposure over eleven seasons. *J Sports Sci* 2020; 38: 238-247
- ³² Gabbett TJ, Ullah S, Finch CF. Identifying risk factors for contact injury in professional rugby league players - application of a frailty model for recurrent injury. *J Sci Med Sport* 2012; 15: 496-504
- ³³ Lauersen JB, Andersen TE, Andersen LB. Strength training as superior, dose-dependent and safe prevention of acute and overuse sports injuries: a systematic review, qualitative analysis and meta-analysis. *Br J Sports Med* 2018; 52: 1557-1563
- ³⁴ Coppack RJ, Etherington J, Wills AK. The effects of exercise for the prevention of overuse anterior knee pain: a randomised controlled trial. *Am J Sports Med* 2011; 39: 940-948
- ³⁵ Askling C, Karlsson J, Thorstensson A. Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. *Scan J Med Sci Sports* 2003; 13: 244-250
- ³⁶ Petersen J, Thorborg K, Neilsen MB et al. Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. *Am J Sports Med* 2011; 39: 2296-2303
- ³⁷ van den Horst N, Smits DW, Petersen J et al. The preventive effect of the Nordic hamstring exercise on hamstring injuries in amateur soccer players: a randomised controlled trial. *Am J Sports Med* 2015; 43: 1316-1323
- ³⁸ Walden M, Atroshi I, Magnussen H et al. Prevention of acute knee injuries in adolescent female football players: cluster randomised controlled trial. *BMJ* 2012; 344: e3042
- ³⁹ Zouita S, Zouita AB, Kebisi W et al. Strength training reduces injury rate in elite young soccer players during one season. *J Strength Cond Res* 2016; 30: 1295-1307
- ⁴⁰ Rugby Football Union. Activate. On the internet: <https://www.englandrugby.com/participation/coaching/activate> accessed 4 May 2020
- ⁴¹ Hislop MD, Stokes KA, Williams S et al. Reducing musculoskeletal injury and concussion risk in schoolboy rugby players with a pre-activity movement control exercise programme: a cluster randomised controlled trial. *Br J Sports Med* 2017; 51: 1140-1146

- ⁴² Attwood MJ, Roberts SP, Trewartha G et al. Efficacy of a movement control injury prevention programme in adult men's community rugby union: a cluster randomised controlled trial. *Br J Sports Med* 2018; 52: 368-374
- ⁴³ Collins CL, Fletcher EN, Fields SK. Neck strength: a protective factor reducing risk for concussion in high school sports. *J Prim Prev* 2014; 35: 309-319
- ⁴⁴ Bengtsson H, Ekstrand J, Waldén M et al. Muscle injury rate in professional football is higher in matches played within 5 days since the previous match: a 14-year prospective study with more than 130 000 match observations. *Br J Sports Med* 2018; 52:1116-1122
- ⁴⁵ Cross MJ, Williams S, Trewartha G et al. The influence of in-season training loads on injury risk in professional rugby union. *Int J Sports Physiol Perform* 2016; 11: 350-355
- ⁴⁶ Williams S, Trewartha G, Kemp SPT et al., How much rugby is too much? A seven-season prospective cohort study of match exposure and injury risk in professional rugby union players. *Sports Med* 2017; 47: 2395-2402.
- ⁴⁷ Williams S, Trewartha G, Kemp S et al. Subsequent injuries and early recurrent diagnoses in elite rugby union players. *Int J Sports Med* 2017; 38: 791-798
- ⁴⁸ Shrier I. Strategic assessment of risk and risk tolerance (StARRT) framework for return-to-play decision-making. *Br J Sports Med* 2015; 49: 1311-1315
- ⁴⁹ Li H, Moreland JJ, Peek-Asa C, Yang J. Preseason Anxiety and Depressive Symptoms and Prospective Injury Risk in Collegiate Athletes. *Am J Sports Med* 2017; 45: 2148-2155
- ⁵⁰ Wall BT, Morton JP, van Loon LJ. Strategies to maintain skeletal muscle mass in the injured athlete: nutritional considerations and exercise mimetics. *Eur J Sport Sci* 2015; 15: 53-62
- ⁵¹ Mitchell JH, Levine BD, McGuire DK. The Dallas bed rest and training study: Revisited after 50 years. *Circulation*. 2019; 140: 1293-1295
- ⁵² Mujika I, Padilla S. Detraining: Loss of training-induced physiological and performance adaptations. Part 1: Short term insufficient training stimulus. *Sport Med* 2000; 30: 79-87
- ⁵³ Dirks ML, Wall BT, van de Valk B et al. One week of bed rest leads to substantial muscle atrophy and induces whole-body insulin resistance in the absence of skeletal muscle lipid accumulation. *Diabetes* 2016; 65: 2862-2875
- ⁵⁴ Dirks ML, Wall BT, Otten B et al. High-fat overfeeding does not exacerbate rapid changes in forearm glucose and fatty acid balance during immobilization. *J Clin Endocrinol Metab* 2020; 105):dgz049.
- ⁵⁵ Kilroe SP, Fulford J, Holwerda AM et al. Short-term muscle disuse induces a rapid and sustained decline in daily myofibrillar protein synthesis rates. *Am J Physiol Endocrinol Metab* 2020; 318: E117-E130

- ⁵⁶ Wall BT, Cruz AM, Otten B et al. The impact of disuse and high-fat overfeeding on forearm muscle amino acid metabolism in humans. *J Clin Endocrinol Metab* 2020; Epub ahead of print
- ⁵⁷ Wall BT, Dirks ML, Snijders T et al. Short-term muscle disuse lowers myofibrillar protein synthesis rates and induces anabolic resistance to protein ingestion. *Am J Physiol Endocrinol Metab* 2016; 310: E137–E147
- ⁵⁸ Wall BT, van Loon LJ. Nutritional strategies to attenuate muscle disuse atrophy. *Nutr Rev* 2013; 71: 195–208
- ⁵⁹ Kilroe SP, Fulford J, Holwerda AM et al. Short-term muscle disuse induces a rapid and sustained decline in daily myofibrillar protein synthesis rates. *Am J Physiol Endocrinol Metab*. 2020;318(2):E117–E130. doi:10.1152/ajpendo.00360.2019
- ⁶⁰ Kilroe SP, Fulford J, Jackman SR et al. Temporal muscle-specific disuse atrophy during one week of leg immobilization. *Med Sci Sports Exerc* 2020; 52: 944–954
- ⁶¹ LeBlanc AD, Schneider VS, Evans HJ et al. Regional changes in muscle mass following 17 weeks of bed rest. *J Appl Physiol* 1992; 73: 2172–2178
- ⁶² Cohen S, Nathan JA, Goldberg AL. Muscle wasting in disease: molecular mechanisms and promising therapies. *Nat Rev Drug Discov*. 2015; 14: 58–74
- ⁶³ Snijders T, Wall BT, Dirks ML et al. Muscle disuse atrophy is not accompanied by changes in skeletal muscle satellite cell content. *Clin Sci (Lond)* 2014; 126: 557–566
- ⁶⁴ Andersen LL, Andersen JL, Magnusson SP, Aagaard P. Neuromuscular adaptations to detraining following resistance training in previously untrained subjects. *Eur J Appl Physiol* 2005; 93: 511–518
- ⁶⁵ Rittweger J, Winwood K, Seynnes O et al. Bone loss from the human distal tibia epiphysis during 24 days of unilateral lower limb suspension. *J Physiol* 2006; 577: 331–337
- ⁶⁶ Dideriksen K, Boesen AP, Reitelseder S et al. Tendon collagen synthesis declines with immobilization in elderly humans: no effect of anti-inflammatory medication. *J Appl Physiol* 2017; 122: 273–282
- ⁶⁷ de Boer MD, Maganaris CN, Seynnes OR et al. Time course of muscular, neural and tendinous adaptations to 23 day unilateral lower-limb suspension in young men. *J Physiol*. 2007; 583: 1079–1091
- ⁶⁸ Kubo K, Ikebukuro T, Maki A et al. Time course of changes in the human Achilles tendon properties and metabolism during training and detraining in vivo. *Eur J Appl Physiol* 2012; 112: 2679–2691
- ⁶⁹ Milsom J, Barreira P, Burgess DJ et al. Case study: Muscle atrophy and hypertrophy in a premier league soccer player during rehabilitation from ACL injury. *Int J Sport Nutr Exerc Metab* 2014; 24: 543–52

- ⁷⁰ *Anderson L, Close GL, Konopinski M et al.* Case Study: Muscle Atrophy, Hypertrophy, and Energy Expenditure of a Premier League Soccer Player During Rehabilitation From Anterior Cruciate Ligament Injury. *Int J Sport Nutr Exerc Metab* 2019; 29: 559-566
- ⁷¹ *García-Pallarés J, Sánchez-Medina L, Carrasco L et al.* Endurance and neuromuscular changes in world-class level kayakers during a periodized training cycle. *Eur J Appl Physiol* 2009; 106: 629-638
- ⁷² *Trappe S, Trappe T, Gallagher P et al.* Human single muscle fibre function with 84 day bed-rest and resistance exercise. *J Physiol* 2004; 557: 501-513
- ⁷³ *Kramer A, Kümmel J, Mulder E et al.* High-Intensity Jump Training Is Tolerated during 60 Days of Bed Rest and Is Very Effective in Preserving Leg Power and Lean Body Mass: An Overview of the Cologne RSL Study. *PLoS One* 2017; 12: e0169793
- ⁷⁴ *Morton RW, Sonne MW, Farias Zuniga A et al.* Muscle fibre activation is unaffected by load and repetition duration when resistance exercise is performed to task failure. *J Physiol* 2019; 597: 4601-4613
- ⁷⁵ *Burd NA, West DW, Staples AW et al.* Low-load high volume resistance exercise stimulates muscle protein synthesis more than high-load low volume resistance exercise in young men. *PLoS One* 2010; 5: e12033
- ⁷⁶ *Mitchell CJ, Churchward-Venne TA, West DW et al.* Resistance exercise load does not determine training-mediated hypertrophic gains in young men. *J Appl Physiol* 2012; 113:71-77
- ⁷⁷ *Lasevicius T, Schoenfeld B, Silva-Batista C et al.* Muscle failure promotes greater muscle hypertrophy in low-load but not in high-load resistance training. *J Strength Cond Res* 2019; Epub ahead of print
- ⁷⁸ *Kubo K, Ikebukuro T, Yata H.* Effects of 4, 8, and 12 Repetition maximum resistance training protocols on muscle volume and strength. *J Strength Cond Res* 2020; Epub ahead of print
- ⁷⁹ *Morton RW, Oikawa SY, Wavell CG et al.* Neither load nor systemic hormones determine resistance training-mediated hypertrophy or strength gains in resistance-trained young men. *J Appl Physiol* 2016; 121: 129-138
- ⁸⁰ *Oxfeldt M, Overgaard K, Hvid LG, Dalgas U.* Effects of plyometric training on jumping, sprint performance, and lower body muscle strength in healthy adults: A systematic review and meta-analyses. *Scand J Med Sci Sports* 2019; 29: 1453-1465
- ⁸¹ *Colliander EB, Tesch PA.* Effects of detraining following short term resistance training on eccentric and concentric muscle strength. *Acta Physiol Scand* 1992; 144: 23-29
- ⁸² *Wilson JM, Marin PJ, Rhea MR et al.,* Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J Strength Cond Res* 2012; 26: 2293-307

- ⁸³ *Hickson RC*. Interference of strength development by simultaneously training for strength and endurance. *Eur J Appl Physiol Occup Physiol* 1980; 45: 255-63
- ⁸⁴ *Mendiguchia J, Conceição F, Edouard P et al*. Sprint versus isolated eccentric training: Comparative effects on hamstring architecture and performance in soccer players. *PLoS One* 2020; 15: e0228283
- ⁸⁵ *Haugen T, Seiler S, Sandbakk Ø, Tønnessen E*. The training and development of elite sprint performance: An integration of scientific and best practice literature. *Sports Med Open* 2019; 5: 44
- ⁸⁶ *Narici MV, Roi GS, Landoni L et al*. Changes in force, cross-sectional area and neural activation during strength training and detraining of the human quadriceps. *Eur J Appl Physiol Occup Physiol* 1989; 59: 310-319
- ⁸⁷ *Brooks SK, Webster RK, Smith LE et al*. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020; 395: 912-920
- ⁸⁸ *Brewer BW, Van Raalte JL, Linder DE*. Athletic identity: Hercules' muscles or Achilles heel? *Int J Sport Psychol* 1993; 24, 237-254.
- ⁸⁹ *Park S, Lavallee D, Tod D*. Athletes' career transition out of sport: A systematic review. *Int Rev Sport Exerc Psychol* 2013; 6: 22-53
- ⁹⁰ *McEwan D, Beauchamp MR*. Teamwork in sport: A theoretical and integrative review. *Int Rev Sport Exerc Psychol* 2014; 7: 229-250
- ⁹¹ *Buchheit M, Morgan W, Wallace J et al*. Physiological, psychometric, and performance effects of the Christmas break in Australian football. *Int J Sports Physiol Perform* 2015; 10: 120-123
- ⁹² *Bompa TO, Buzzichelli C*. Periodization: Theory and methodology of training. Champaign, IL: Human Kinetics, 2018.
- ⁹³ *Grout H, Long G*. Improving teaching and learning in physical education. Maidenhead: McGraw-Hill Education, 2009
- ⁹⁴ *Mujika I, Halson S, Burke LM, et al*. An integrated, multifactorial approach to periodization for optimal performance in individual and team sports. *Int J Sports Physiol Perform* 2018 13; 538-561
- ⁹⁵ *Farrow D, Robertson S*. Development of a skill acquisition periodisation framework for high-performance sport. *Sports Med* 2017; 47: 1043-1054
- ⁹⁶ *Martin KA, Moritz SE, Hall CR*. Imagery use in sport: A literature review and applied model. *Sport Psychol* 1999; 13: 245-268
- ⁹⁷ *Bruton AM, Mellalieu SD, Shearer DA*. Observation interventions as a means to manipulate collective efficacy in groups. *J Sport Exerc Psychol* 2014; 36: 27-39

- ⁹⁸ *Mellalieu, S. D., Sellars, P. A., Wagstaff, C. R. D., Arnold, R., & Neil, R. (2020). Understanding the psychological load and successful coping of rugby union players in the English professional game. Cardiff Metropolitan University, UK.*
- ⁹⁹ *Eccles D, Balk Y, Gretton T, Harris N. "The Forgotten Session": Advancing research and practice concerning the psychology of rest in athletes. J Appl Sport Psychol 2020; Epub Ahead of print*
- ¹⁰⁰ *Tedeschi RG, Shakespeare-Finch J, Taku K, Calhoun LG. Posttraumatic growth: Theory, research, and applications. Routledge, 2018*
- ¹⁰¹ *Howells K, Sarkar M, Fletcher D. Can athletes benefit from difficulty? A systematic review of growth following adversity in competitive sport. Prog Brain Res 2017; 16: 37–48.*
- ¹⁰² *Wadey R, Day M, Howells K. Growth following adversity in sport: A mechanism to positive change in sport. Routledge, 2020*
- ¹⁰³ *Anderson L, Naughton RJ, Close GL et al. Daily distribution of macronutrient intakes of professional soccer players from the English Premier League. Int J Sport Nutr Exerc Metab 2017; 28: 1-18*
- ¹⁰⁴ *Anderson L, Orme P, Naughton RJ et al. Energy intake and expenditure of professional soccer players of the English Premier League: evidence of carbohydrate periodization. Int J Sport Nutr Exerc Metab 2017; 27: 228-238*
- ¹⁰⁵ *Anderson L, Close GL, Konopinski M et al. Case study: Muscle atrophy, hypertrophy and energy expenditure of a Premier League soccer player during rehabilitation from ACL injury. Int J Sport Nutr Exerc Metab 2019; 29: 559-566*
- ¹⁰⁶ *Morehen JC, Bradley WJ, Clarke J et al. The assessment of total energy expenditure during a 14-day in-season period of professional rugby league players using the doubly labelled water method. Int J Sport Nutr Exerc Metab 2016; 26: 464-472*
- ¹⁰⁷ *Smith D, King RFGJ, Duckworth LC et al. Energy expenditure of rugby players during a 14-day in-season period, measured using doubly labelled water. Eur J Appl Physiol 2018; 118: 647-656*
- ¹⁰⁸ *Costello N, Deighton K, Preston T et al. Are professional young rugby league players eating enough? Energy intake, expenditure and balance during pre-season. Eur J Sport Sci 2019; 19: 123-132*
- ¹⁰⁹ *Costello N, Deighton K, Preston T et al. (2018) Collisions activity during training increase total energy expenditure measured via doubly labelled water. Eur J Appl Physiol 2018; 118: 1169-1177*
- ¹¹⁰ *Olsen RH, Krogh-Madsen R, Thomsen C et al. Metabolic responses to reduced daily steps in healthy nonexercising men. JAMA 2008; 299: 1261-1263*

- 1212 ¹¹¹ *Knudsen SH, Hansen LS, Pedersen M et al.* Changes in insulin sensitivity precede changes
1213 in body composition during 14 days of step reduction combined with overfeeding in healthy
1214 young men. *J Appl Physiol* 2012; 113: 7-15
1215
- 1216 ¹¹² *English KL, Mettler JA, Ellison JB et al.* Leucine partially protects muscle mass and
1217 function during bed rest in middle-aged adults. *Am J Clin Nutr* 2016; 103: 465-473
1218
- 1219 ¹¹³ *Kortebein P, Symons TB, Ferrando A et al.* Functional impact of 10 days of bed rest in
1220 healthy older adults. *J Gerontol A Biol Sci Med Sci* 2008; 63: 1076-1081
1221
- 1222 ¹¹⁴ *Biolo G, Ciocchi B, Stulle M et al.* Calorie restriction accelerates the catabolism of lean
1223 body mass during 2 wk of bed rest. *Am J Clin Nutr*, 2007; 86: 366-372
1224
- 1225 ¹¹⁵ *Paddon-Jones D, Sheffield-Moore M, Urban RJ et al.* Essential amino acid and
1226 carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days
1227 bedrest. *J Clin Endocrinol Metab* 2004; 89: 4351-4358
1228
- 1229 ¹¹⁶ *Churchward-Venne TA, Burd NA et al.* Supplementation of a suboptimal dose with leucine
1230 or essential amino acids: Effects on myofibrillar protein synthesis at rest and following
1231 resistance exercise in men. *J Physiol* 2012; 590: 2751-2765
1232
- 1233 ¹¹⁷ *Morton RW, Murphy KT, McKellar SR, et al.* A systematic review, meta-analysis and
1234 meta-regression of the effect of protein supplementation on resistance training-induced gains
1235 in muscle mass and strength in healthy adults. *B J Sports Med* 2018; 52: 376-384
1236
- 1237 ¹¹⁸ *Impey SG, Hearris MA, Hammond KM et al.* Fuel the work required: A theoretical
1238 framework for carbohydrate periodization and the glycogen threshold hypothesis. *Sports Med*
1239 2018; 48: 1031-1048.
1240
- 1241 ¹¹⁹ *Nieman DC, Hazels Mitmesser S.* Potential impact of nutrition on immune system
1242 recovery from heavy exertion: A metabolomics perspective. *Nutrients* 2017; 9: 513-536
1243
- 1244 ¹²⁰ *Walsh NP.* Nutrition and athlete immune health: New perspectives on an old paradigm.
1245 *Sports Med* 2019; 49: S153-S168
1246
- 1247 ¹²¹ *Williams NC, Killer SC, Svendsen IS, Wyn Jones A.* Immune nutrition and exercise:
1248 Narrative review and practical recommendations. *Eur J Sport Sci* 2018; 19: 49-61
1249
- 1250 ¹²² *Hespeel P, Op't Eijnde B, Van Leemputte M et al.* Oral creatine supplementation facilitates
1251 the rehabilitation of disuse atrophy and alters the expression of muscle myogenic factors in
1252 humans. *J Phys* 2001; 536: 625-633
1253
- 1254 ¹²³ *Johnston AP, Burke DG, MacNeil LG, Candow DG.* Effect of creatine supplementation
1255 during cast-induced immobilization on the preservation of muscle mass, strength, and
1256 endurance. *J Strength Cond Res* 2009; 23: 116-120
1257
- 1258 ¹²⁴ *Nova, E., Varela, P., Lopez-Vidriero, I., et al.* A one-year follow-up study in anorexia
1259 nervosa: Dietary pattern and anthropometrical evolution. *Eur J Clin Nutr* 2001; 55: 574-554
1260
- 1261 ¹²⁵ *Woodward, B.* Protein, calories, and immune function. *Nutr Rev* 1998; 56: S84-S92.

- ¹²⁶ Hume R, Weyers E. Changes in leucocyte ascorbic acid during the common cold. *Scott Med J* 1973; 18: 3-7.
- ¹²⁷ He CS, Handzlik M, Fraser WD, et al. Influence of vitamin D status on respiratory infection incidence and immune function during 4 months of winter training in endurance sport athletes. *Exercise Immunology Review* 2013; 19: 86-101
- ¹²⁸ Martineau AR, Jolliffe DA, Hooper RL et al. Vitamin D supplementation to prevent acute respiratory tract infections: Systematic review and meta-analysis of individual participant data. *Br Med J* 2017; 356: i6583
- ¹²⁹ Copley C, Cox AJ, Pyne DB et al. Upper respiratory symptoms, gut health and mucosal immunity in athletes. *Sports Med* 2018; 48: 65-77
- ¹³⁰ Pyne DB, West NP, Cox AJ, Cripps AW. Probiotics supplementation for athletes: Clinical and physiological effects. *Eur J Sport Sci* 2015; 15: 63-72
- ¹³¹ Ayres JS, Sneider DS. Tolerance of infections. *Annu Rev Immunol* 2012; 30, 271-294.
- ¹³² Hendricks S, Till K, den Hollander S et al., Consensus on a video analysis framework of descriptors and definitions by the Rugby Union Video Analysis Consensus group. *Br J Sport Med* 2020; Epub Ahead of Print
- ¹³³ Sclafani MP, Davis CC. Return to play progression for rugby following injury to the lower extremity: a clinical commentary and review of the literature. *Int J Sports Phys Ther* 2016; 11: 302-32
- ¹³⁴ Jiménez-Reyes P, Cross M, Ross A et al. Changes in mechanical properties of sprinting during repeated sprint in elite rugby sevens athletes. *Eur J Sport Sci* 2019; 19: 585-594
- ¹³⁵ Buckthorpe M, Wright S, Bruce-Low S et al. Recommendations for hamstring injury prevention in elite football: translating research into practice. *Br J Sports Med*. 2019; 53: 449-456
- ¹³⁶ Haugen T, Danielsen J, McGhie D et al. Kinematic asymmetry in the stride cycle is not associated with performance and injuries in athletic sprinters. *Scand J Med Sci Sports*. 2018; 28: 1001–1008
- ¹³⁷ Haugen T, Tønnessen E, Hisdal J, Seiler S. The role and development of sprinting speed in soccer. *Int J Sports Physiol Perform* 2014; 9: 432–441
- ¹³⁸ Weaving D, Marshall P, Earle K et al. Quantifying the external and internal loads of professional rugby league training modes: consideration for concurrent field-based training prescription. *J Strength Cond Res* 2017; Epub ahead of print
- ¹³⁹ Petrakos G, Morin JB, Egan B. Resisted sled sprint training to improve sprint performance: A systematic review. *Sports Med* 2016; 46: 381-400

- ¹⁴⁰ Francis C. Structure of training for speed (ebook). [https://www.amazon.com/ Structure-Training-Charlie-Francis-Concepts-ebook/dp/B00BG9F8UG](https://www.amazon.com/Structure-Training-Charlie-Francis-Concepts-ebook/dp/B00BG9F8UG).
- ¹⁴¹ Green BS, Blake C, Caulfield BM. A comparison of cutting technique performance in rugby union players. *J Strength Cond Res* 2011; 25: 2668-2680
- ¹⁴² Spiteri T, Hart NH, Nimphius S. Offensive and defensive agility: a sex comparison of lower body kinematics and ground reaction forces. *J Appl Biomech* 2014; 30: 514-520
- ¹⁴³ Hendricks S, Till K, Brown J, Jones B. Rugby Union needs a contact skill-training programme, *Brit J Sports Med* 2017; 51: 829-830
- ¹⁴⁴ Burger N, Lambert MI, Viljoen W et al. Tackle technique and tackle-related injuries in high-level South African Rugby Union under-18 players: real-match video analysis. *Br J Sports Med* 2016; 50: 932-938
- ¹⁴⁵ Burger N, Lambert MI, Viljoen W et al. Mechanisms and factors associated with tackle-related injuries in South African youth rugby union players. *Am J Sports Med* 2017; 45: 278-285
- ¹⁴⁶ Tierney GJ, Denvir K, Farrell G, Simms CK. Does player time-in-game affect tackle technique in elite level rugby union? *J Sci Med Sport* 2018; 21: 221-225
- ¹⁴⁷ Davidow D, Redman M, Lambert M et al. The effect of physical fatigue on tackling technique in rugby union. *J Sci Med Sport* 2020; Epub Ahead of print
- ¹⁴⁸ Hendricks S, Till K, Oliver J et al. A technical skill training framework and skill load measurement for the rugby union tackle. *Strength and Cond J* 2018; 40: 44-49
- ¹⁴⁹ Hendricks S, Till K, Oliver J et al. Rating of perceived challenge as a measure of internal load for technical skill performance. *Brit J Sports Med* 2019; 53: 611-613
- ¹⁵⁰ Issurin V. Block Periodization: Breakthrough in Sports Training. Ultimate athlete concepts, 2008
- ¹⁵¹ Kanehisa H, Miyashita M. Specificity of velocity in strength training. *Eur J Appl Physiol Occup Physiol* 1983; 52: 104-106
- ¹⁵² Behm DG, Sale DG. Velocity specificity of resistance training. *Sport Med* 1993; 15(6):374-388.
- ¹⁵³ Zamparo P, Minetti A, Di Prampero P. Interplay among the changes of muscle strength, cross-sectional area and maximal explosive power: theory and facts. *Eur J Appl Physiol* 2002; 88: 193-202.
- ¹⁵⁴ Snijders T, Aussieker T, Holwerda A, et al. The concept of skeletal muscle memory: Evidence from animal and human studies. *Acta Physiol (Oxf)* 2020; Epub ahead of print

- ¹⁵⁵ *Staron RS, Leonardi MJ, Karapondo DL, et al.* Strength and skeletal muscle adaptations in heavy-resistance-trained women after detraining and retraining. *J Appl Physiol* 1991; 70: 631-640
- ¹⁵⁶ *The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team* The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Chin J Epidemiol* 2020; 41: Epub ahead of print. DOI: 10.3760/cma.j.issn.0254-6450.2020.02.003
- ¹⁵⁷ *Shi S, Qin M, Shen B et al.* Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol* 2020; 25: Epub ahead of print
- ¹⁵⁸ *Liu P, Blet A, Smyth D, Hongliang Li.* The Science Underlying COVID-19: Implications for the Cardiovascular System. *Circulation* 2020; 25: Epub ahead of print
- ¹⁵⁹ *Eichner ER.* Infection, immunity, and exercise. *Phys Sportsmed* 1993; 21: 125-135
- ¹⁶⁰ *Kim IC, Kim JY, Kim HA, Han S.* COVID-19-related myocarditis in a 21-year-old female patient. *Eur Heart J* 2020; 13; Epub ahead of print
- ¹⁶¹ *Kenney K, Landau ME, Gonzalez RS et al.* Serum creatine kinase after exercise: drawing the line between physiological response and exertional rhabdomyolysis. *Muscle Nerve* 2012; 45: 356-362
- ¹⁶² *Meeusen R, Duclos M, Foster C et al.* Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. *Med Sci Sports Exerc* 2013; 45: 186-205
- ¹⁶³ *Quarrie KL, Raftery M, Blackie J et al.* Managing player load in professional rugby union: a review of current knowledge and practices. *Br J Sports Med* 2017; 51: 421-427
- ¹⁶⁴ *Foster C, Florhaug J, Franklin et al.* A new approach to monitoring exercise training. *J Strength Cond Res* 2001; 15: 109-115
- ¹⁶⁵ *Franchini M, Ferraresi M, Pretruo A et al.* Is a retrospective RPE appropriate in soccer? Response shift and recall bias. *Sci Med Football* 2017; 1: 53-59
- ¹⁶⁶ *Scantlebury S, Till K, Sawczuk T et al.* Validity of Retrospective Session Rating of Perceived Exertion to Quantify Training Load in youth Athletes. *J Strength Cond Res* 2018; 32: 1975-1980
- ¹⁶⁷ *Phibbs PJ, Roe G, Jones B et al.* Validity of Daily and Weekly Self-Reported Training Load Measures in Adolescent Athletes. *J Strength Cond Res* 2017; 31: 1121-1126
- ¹⁶⁸ *Harriss DJ, Macsween A, Atkinson G.* Ethical Standards in Sport and Exercise Science Research: 2020 Update. *Int J Sports Med* 2019; 40: 813-817

Figure legends

Figure 1. Incidence (A: injuries per 1000 hours) and burden (B: days lost per 1000 hours) of training injuries during the pre-season and early competition period in the English Premiership (2014-15 to 2018-19 seasons)

Figure 2. Training considerations following return to play after the period of restricted training due to COVID-19. Reintroduction of group training will require progressions and structure of training to be developed with reference to risk of COVID-19 transmission.

Figure 3. Return to play risk stratification for athletes following COVID-19 symptoms.

Figure 4. A summary of 'at risk' athletes following modified training due to COVID-19.

Table legends

Table 1. Energy expenditures of professional and elite male rugby players during various stages of the season, measured via doubly labelled water (DLW).

Table 2. Most effective intensity and volume for strength and power in collision sports and the expected percentage change in maximal strength and power per training session conducted (data from McMaster et al., [3]).

Table 3. The impact of training frequency weekly strength and power changes (data from McMaster et al., [3]).

Table 4. Rate of changes in strength and power markers in a tier 1 international rugby union team over 5-weeks (n=26 players).

Table 5. Changes in strength markers in a professional rugby union team over 4-weeks (n=35 players).

Table 6. Challenges and practical recommendations for sports during and following COVID-19.